



AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings of claims in the application:

LISTING OF CLAIMS:

1-23. (canceled)

24. (currently amended) ~~Compounds A~~ A compound of the following general formula (I):



wherein R_1 and R_2 are fatty acyl groups.

25. (currently amended) ~~Compounds~~ The compound according to claim 24, wherein R_1 and R_2 independently from each other represent a fatty acyl group containing from 16 to 60 carbon atoms, and ~~more particularly~~ wherein R_1 and R_2 are selected from the group comprising:


- $\text{—C(=O)—(CH}_2\text{)}_{14}\text{—CH}_3$ (palmitic acyl),
- $\text{—C(=O)—(CH}_2\text{)}_{16}\text{—CH}_3$ (stearic acyl),
- $\text{—C(=O)—(CH}_2\text{)}_{18}\text{—CH}_3$ (arachidic acyl),
- $\text{—C(=O)—(CH}_2\text{)}_{20}\text{—CH}_3$ (docosanoic acyl),
- $\text{—C(=O)—(CH}_2\text{)}_{22}\text{—CH}_3$ (tetracosanoic acyl),
- $\text{—C(=O)—CH(CH}_3\text{)—[CH}_2\text{—CH(CH}_3\text{)]}_n\text{—CH(OH)—[CH}_2\text{]}_m\text{—CH}_3$ (hydroxyphthioceranoic acyl),

wherein m is 14 or 16 and n is an integer from 2 to 10,

- $\text{—C(=O)—CH(CH}_3\text{)—[CH}_2\text{—CH(CH}_3\text{)]}_n\text{—CH}_2\text{—[CH}_2\text{]}_m\text{—CH}_3$ (phthioceranoic acyl),

wherein m is 14 or 16 and n is an integer from 2 to 10,

26. (currently amended) ~~Compounds~~ The compound according to claim 24, wherein R_1 and R_2 are selected from the group ~~comprising~~ consisting of palmitic acyl and stearic acyl, ~~namely compounds of following formulae:~~ and having the formula:





27. (currently amended) ~~Compounds~~ The compound according to claim 24, wherein at least one of R₁ and R₂ represents a hydroxyphthioceranoic acyl group.

28. (currently amended) ~~Compounds~~ The compound according to claim 24, wherein R₁ or R₂ represents a hydroxyphthioceranoic acyl group.



29. (currently amended) ~~Compounds~~ The compound according to claim 24, wherein:

- R_1 represents a hydroxyphthioceranoic acyl group, and R_2 represents a palmitic acyl group or a stearic acyl group, namely compounds of following formula (II):




wherein p is 14 or 16, m is 14 or 16 and n is an integer from 2 to 10, or


- R_2 represents a hydroxyphthioceranoic acyl group, and R_1 represents a palmitic acyl group or a stearic acyl group, namely compounds of following formula (III):



wherein p is 14 or 16, m is 14 or 16 and n is an integer from 2 to 10.

30.(currently amended) ~~Compounds~~ The compound according to claim 24, of following formula II,





wherein the compound is selected from the group consisting of:

- $n = 2, m = 14$ and $p = 14$ (II.1) ;
- $n = 2, m = 14$ and $p = 16$ (II.2) ;
- $n = 2, m = 16$ and $p = 14$ (II.3) ;
- $n = 2, m = 16$ and $p = 16$ (II.4) ;

- $n = 3, m = 14$ and $p = 14$ (II.5) ;
- $n = 3, m = 14$ and $p = 16$ (II.6) ;
- $n = 3, m = 16$ and $p = 14$ (II.7) ;
- $n = 3, m = 16$ and $p = 16$ (II.8) ;

- $n = 4, m = 14$ and $p = 14$ (II.9) ;
- $n = 4, m = 14$ and $p = 16$ (II.10) ;
- $n = 4, m = 16$ and $p = 14$ (II.11) ;
- $n = 4, m = 16$ and $p = 16$ (II.12) ;

- $n = 5, m = 14$ and $p = 14$ (II.13) ;
- $n = 5, m = 14$ and $p = 16$ (II.14) ;
- $n = 5, m = 16$ and $p = 14$ (II.15) ;
- $n = 5, m = 16$ and $p = 16$ (II.16) ;

- $n = 6, m = 14$ and $p = 14$ (II.17) ;
- $n = 6, m = 14$ and $p = 16$ (II.18) ;

- $n = 6, m = 16$ and $p = 14$ (II.19) ;
- $n = 6, m = 16$ and $p = 16$ (II.20) ;


- $n = 7, m = 14$ and $p = 14$ (II.21) ;
- $n = 7, m = 14$ and $p = 16$ (II.22) ;
- $n = 7, m = 16$ and $p = 14$ (II.23) ;
- $n = 7, m = 16$ and $p = 16$ (II.24) ;


- $n = 8, m = 14$ and $p = 14$ (II.25) ;
- $n = 8, m = 14$ and $p = 16$ (II.26) ;
- $n = 8, m = 16$ and $p = 14$ (II.27) ;
- $n = 8, m = 16$ and $p = 16$ (II.28) ;

- $n = 9, m = 14$ and $p = 14$ (II.29) ;
- $n = 9, m = 14$ and $p = 16$ (II.30) ;
- $n = 9, m = 16$ and $p = 14$ (II.31) ;
- $n = 9, m = 16$ and $p = 16$ (II.32) ;

- $n = 10, m = 14$ and $p = 14$ (II.33) ;
- $n = 10, m = 14$ and $p = 16$ (II.34) ;
- $n = 10, m = 16$ and $p = 14$ (II.35) ; and
- $n = 10, m = 16$ and $p = 16$ (II.36) ;

or of following formula III,





wherein[[:]] the compound is selected from the group consisting of

- $n = 2, m = 14$ and $p = 14$ (III.1) ;
- $n = 2, m = 14$ and $p = 16$ (III.2) ;
- $n = 2, m = 16$ and $p = 14$ (III.3) ;
- $n = 2, m = 16$ and $p = 16$ (III.4) ;

- $n = 3, m = 14$ and $p = 14$ (III.5) ;
- $n = 3, m = 14$ and $p = 16$ (III.6) ;
- $n = 3, m = 16$ and $p = 14$ (III.7) ;
- $n = 3, m = 16$ and $p = 16$ (III.8) ;

- $n = 4, m = 14$ and $p = 14$ (III.9) ;
- $n = 4, m = 14$ and $p = 16$ (III.10) ;
- $n = 4, m = 16$ and $p = 14$ (III.11) ;
- $n = 4, m = 16$ and $p = 16$ (III.12) ;

- $n = 5, m = 14$ and $p = 14$ (III.13) ;
- $n = 5, m = 14$ and $p = 16$ (III.14) ;
- $n = 5, m = 16$ and $p = 14$ (III.15) ;
- $n = 5, m = 16$ and $p = 16$ (III.16) ;

- $n = 6, m = 14$ and $p = 14$ (III.17) ;

- $n = 6, m = 14$ and $p = 16$ (III.18) ;
- $n = 6, m = 16$ and $p = 14$ (III.19) ;
- $n = 6, m = 16$ and $p = 16$ (III.20) ;




- $n = 7, m = 14$ and $p = 14$ (III.21) ;
- $n = 7, m = 14$ and $p = 16$ (III.22) ;
- $n = 7, m = 16$ and $p = 14$ (III.23) ;
- $n = 7, m = 16$ and $p = 16$ (III.24) ;


- $n = 8, m = 14$ and $p = 14$ (III.25) ;
- $n = 8, m = 14$ and $p = 16$ (III.26) ;
- $n = 8, m = 16$ and $p = 14$ (III.27) ;
- $n = 8, m = 16$ and $p = 16$ (III.28) ;

- $n = 9, m = 14$ and $p = 14$ (III.29) ;
- $n = 9, m = 14$ and $p = 16$ (III.30) ;
- $n = 9, m = 16$ and $p = 14$ (III.31) ;
- $n = 9, m = 16$ and $p = 16$ (III.32) ;


- $n = 10, m = 14$ and $p = 14$ (III.33) ;
- $n = 10, m = 14$ and $p = 16$ (III.34) ;
- $n = 10, m = 16$ and $p = 14$ (III.35) ; and
- $n = 10, m = 16$ and $p = 16$ (III.36) [[;]].

31.(currently amended) Compounds according to claim 24, of following formulae:







11.21




III.22




III.22




III.23



III.23





II.23



32. (previously presented) A composition comprising at least two different compounds of formula I such as defined in claim 24.

33. (currently amended) A composition according to claim 32, characterized in that it comprises a mixture of compounds selected from the compounds of following formula (II):







11




wherein p is 14 or 16, m is 14 or 16 and n is an integer from 2 to 10, or





from the compounds of following formula (III):








wherein p is 14 or 16, m is 14 or 16 and n is an integer from 2 to 10.

34. (currently amended) A composition according to claim 32, characterized in that it comprises a mixture of compounds selected from the following compounds:









- 35.(currently amended) [[A]] The composition according to claim 34, wherein the compounds represent from about 20% to about 100%, more particularly about 30%, of the total amount of compounds of formula I of said composition.
- 36.(previously presented) A pharmaceutical composition comprising at least one compound of claim 24, in association with a pharmaceutically acceptable vehicle.
- 37.(currently amended) [[A]] The pharmaceutical composition according to claim 36, characterized in that it is presented in a form intended for administration by oral or injectable route.
- 38.(previously presented) A pharmaceutical composition according to claim 36, characterized in that it comprises one or more other products useful for the treatment or the prophylaxis of tuberculosis, such as BCG or mycobacterial proteins.
- 39.(currently amended) ~~Products~~ A product comprising:
- at least one compound of claim 24,
 - and at least one other product useful for the treatment or the prophylaxis of tuberculosis, ~~such as BCG or mycobacterial proteins,~~

as a combined preparation for simultaneous, separate or sequential use in the treatment or the prophylaxis of tuberculosis.

40.(currently amended) ~~Method~~ A method for the treatment or the prophylaxis of tuberculosis, comprising the administration of a therapeutically effective amount to a patient of at least one compound of claim 24.

41.(currently amended) ~~Method~~ A method of activating immune reaction, ~~and more particularly for activating inflammatory reaction,~~ comprising the administration of a therapeutically effective amount to a patient of at least one compound of claim 24.

42.(currently amended) ~~Method~~ A method of inducing the activation of T lymphocytes, comprising the administration of a therapeutically effective amount to a patient of at least one compound of claim 24.

43.(currently amended) ~~Method~~ A method of inducing the production of IFN- γ , TNF- α , IL-4 or granulysin, comprising the administration of a therapeutically effective amount to a patient of at least one compound of claim 24.

44.(currently amended) A process for generating T cell clones, characterized in that ~~[[it]]~~ said process comprises the following stages:

- incubating antigen presenting cells (APCs), ~~notably dendritic cells,~~ with a *Mycobacterium tuberculosis* envelope preparation substantially devoid of proteins, to obtain non-protein envelope antigen loaded APCs,

- contacting peripheral blood mononuclear cells with the envelope antigen loaded APCs to obtain proliferating T cells,
- cloning proliferating T cells by limiting dilution and selecting the clones releasing a molecule selected from the group comprising IFN- γ , TNF- α , granulysin or IL-4 when contacted by envelope antigen loaded APCs to obtain T cell clones.

45.(previously presented) T cell clones such as obtained by the process according to claim 44.

46.(currently amended) A process for screening products, such as sulfoglycolipids extracted from *Mycobacterium tuberculosis*, characterised in that ~~[[it]]~~ said process comprises the following stages: .

- contacting dendritic cells loaded with the product to screen, notably sulfoglycolipids extracted from *Mycobacterium tuberculosis*, with T cell clones according to claim 45,
- detecting a molecule selected from the group ~~comprising~~ consisting of IFN- γ , TNF- α , granulysin ~~[[or]]~~ and IL-4, released by the T cell clones.

47.(currently amended) A process for the extraction of compounds of claim 24, from *Mycobacterium tuberculosis*, characterized in that ~~[[it]]~~ said process comprises the following stages:

- treatment of *M. tuberculosis* bacteria with a mixture of methanol and chloroform to obtain a chloroform/methanol extract,

- concentration of the chloroform/methanol extract followed by its partition between a chloroform phase and an aqueous phase,
- taking of the chloroform phase and evaporation of most of the chloroform, followed by addition of acetone thereto to obtain a precipitate and a soluble acetone phase,
- taking of the soluble acetone phase followed by [[its]] concentration, and application of the concentrated soluble acetone phase on a silicic acid column irrigated with mixtures of methanol and chloroform,
- elution of a fraction from the above-mentioned silicic acid column by a mixture of chloroform and approximately 20% methanol,
- if necessary purification of the fraction eluted from the silicic acid column to obtain different preparations respectively containing substantially only one compound.